

Marginal Deformations In the Open Bosonic String Field Theory for N D0-branes

J. Klusoň *

Institute of Theoretical Physics, University of Stockholm, SCFAB

SE- 106 91 Stockholm, Sweden

and

Institutionen för teoretisk fysik

BOX 803, SE- 751 08 Uppsala, Sweden

E-mail: josef.kluson@teorfys.uu.se

ABSTRACT: In this short note we give an example of the exact solution of the open bosonic string field theory defined on the background of N coincided D0-branes. This solution leads to the change of the original background to the background where D0-branes are localised in general positions.

KEYWORDS: String field theory.

*On leave from Masaryk University, Brno

Contents

1. Introduction	1
2. Open bosonic string field theory for N D0-branes	2
3. Finite deformation	11
4. Conclusion	13

1. Introduction

In our recent paper [14] we gave examples of two exact solutions of the open bosonic string field theory [4] that correspond to the marginal deformation of the original background configuration of D0-brane sitting in the origin. These two examples served as a demonstration of the efficiency of our method developed in [13]. We have seen that the solutions given in [14] did not change the form of the BRST operator and hence the spectrum of the string states around the new background is the same. As was argued in that paper this result is a trivial consequence of the homogeneity and isotropy of the space where D0-brane lives. We have also argued that in the case of N D0-branes the situation will not be so simple. The analysis of this configuration will be performed in this paper where we find exact solution of the open bosonic string field theory that corresponds to the marginal deformation of the original configuration. We will see that when we perform expansion around the new background configuration we obtain the string field theory action for the fluctuation field that has the same form as the original one however the new BRST operator is matrix valued as a consequence of the nonabelian nature of the initial configuration. This result supports the proposal given in [15], however this claim certainly deserves much more detailed study and we will postpone it for the future. Then we will argue, following our general discussion given in [13], that we can perform the field redefinition that maps the new BRST operator to the original one and the fluctuation field that arises from the expansion of the string field around the classical solution of the string field theory equation of motion to the fluctuation field with nonzero winding number. More precisely, we will see that the state of the string joining i -Th and j -Th D0-branes has the winding number proportional to the distance of these two D0-branes.

Since this paper is closely related to [13, 14] we will not discuss the general construction presented in [13] and the abelian solution given in [14], where more details and more detailed list of references can be found. However it is a pleasant duty for us mention some of the papers [5, 6, 7, 8, 9] that discuss related problems from different point of view.

In the conclusion we outline our results and suggest possible extension of our work.

2. Open bosonic string field theory for N D0-branes

In this section we present nonabelian generalization of our result given in [14]. For more details and motivation, see that paper.

Let us consider system of coincident N D0-branes localized in the origin of the target space. As was argued in [4] such a system is described with the standard open bosonic string field theory where now string field will carry Chan-Paton (CP) indexes so that we can write

$$S = - \sum_{i,j,k} \int \frac{1}{2} \Phi_{ij} \star Q(\Phi_{ji}) + \frac{1}{3} \Phi_{ij} \star \Phi_{jk} \star \Phi_{ki} , \quad (2.1)$$

where Φ_{ij} is a ghost number one string field and the summation index goes from 1 to N and where Q is the BRST operator of open string that obeys Dirichlet boundary conditions in $x^I, I = 1, \dots, 25$ directions (For more details about string field theory, see very good reviews [1, 2, 3]). From (2.1) we obtain the equation of motion

$$Q\Phi_{ij} + \sum_k \Phi_{ik} \star \Phi_{kj} = 0 . \quad (2.2)$$

The generalization of the solution given in [14] to the nonabelian case is straightforward

$$\Phi_{0ij} = \sum_k (e^{-\Psi_0})_{ik} \star Q(e^{\Psi_0})_{kj} . \quad (2.3)$$

It is easy to see that Φ_{0ij} with any ghost number zero field Ψ_{0ij} is solution of the equation of motion (2.2) since we have

$$Q(\Phi_0)_{ij} = Q \left(\sum_k (e^{-\Psi_0})_{ik} \star Q(e^{\Psi_0})_{kj} \right) = - \sum_{k,l,m} (e^{-\Psi_0})_{ik} \star Q(e^{\Psi_0})_{kl} \star (e^{-\Psi_0})_{lm} \star Q(e^{-\Psi_0})_{mj} \quad (2.4)$$

and

$$\sum_k (\Phi_0)_{ik} \star (\Phi_0)_{kj} = \sum_{k,m,l} (e^{-\Psi_0})_{ik} \star Q(e^{\Psi_0})_{kl} \star (e^{-\Psi_0})_{lm} \star Q(e^{\Psi_0})_{mj} . \quad (2.5)$$

As in case of single D0-brane [14] we propose that the field Ψ_{0ij} has the form

$$(\Psi_0)_{ij} = \hat{K}_L(\hat{I}, F)_{ij} , \quad (2.6)$$

where we define the generalization of the identity field of the string star algebra to the nonabelian case as

$$\hat{I}_{ij} = \mathcal{I} \otimes \delta_{ij} . \quad (2.7)$$

This field has the basic property of the identity element of the string field algebra

$$\sum_k \Psi_{ik} \star \hat{I}_{kj} = \Psi_{ij} \star \mathcal{I} = \Psi_{ij} , \sum_k \hat{I}_{ik} \star \hat{I}_{kj} = \hat{I}_{ij} . \quad (2.8)$$

In (2.6) we have also introduced the ghost number zero operator $\hat{K}$. Following [13] we will study the fluctuation around the solution Φ_{0ij} . We write general string field as follows

$$\Phi_{ij} = \Phi_{0ij} + \Psi_{ij} \quad (2.9)$$

and insert it into (2.1). According to the well known procedure (For recent discussion, see [10].) we obtain the action for the field Ψ_{ij} which has the same form as the original one (2.1) however with the new BRST operator

$$\tilde{Q}(X)_{ij} = \sum_k \delta_{ik} Q(X)_{kj} + \Phi_{0ik} \star X_{kj} - (-1)^{|X|} X_{ik} \star \Phi_{0kj} . \quad (2.10)$$

According to [13] this new BRST operator $\tilde{Q}$ has a form

$$\tilde{Q}(X)_{ij} = e^{-\hat{K}} \left(Q \left(e^{\hat{K}}(X) \right) \right)_{ij} . \quad (2.11)$$

In order the string field theory for fluctuation field Ψ_{ij} to be well defined we must show that the BRST operator (2.11) obeys all axioms of the open bosonic string field theory [1, 2, 3].

Let us start with the proof of the nilpotence of (2.30)²

$$\begin{aligned} \tilde{Q}^2(X)_{ij} &= \tilde{Q} \left(Q(X)_{ij} + \Phi_{0ik} \star X_{kj} - (-1)^{|X|} X_{ik} \star \Phi_{0kj} \right) = \\ &= Q \left(Q(X)_{ij} + \Phi_{0ik} \star X_{kj} - (-1)^{|X|} X_{ik} \star \Phi_{0kj} \right) + \\ &\quad + \Phi_{0ik} \star \left(Q(X)_{kj} + \Phi_{0kl} \star X_{lj} - (-1)^{|X|} X_{kl} \star \Phi_{0lj} \right) - \\ &\quad - (-1)^{|X|+1} \left(Q(X)_{ik} + \Phi_{0ik} \star X_{kl} - (-1)^{|X|} X_{ik} \star \Phi_{0kl} \right) \star \Phi_{0lj} = \\ &= (Q(\Phi_{0ik}) + \Phi_{0il} \star \Phi_{0lk}) \star X_{kj} - (-1)^{2|X|} X_{ik} \star (Q(\Phi_{0kj}) + \Phi_{0kl} \star \Phi_{0lj}) = 0 \end{aligned} \quad (2.12)$$

since Φ_{0ij} obeys (2.2). Derivation property can be seen very easily since we have

$$\begin{aligned} \tilde{Q}(X \star Y)_{ij} &= Q(X)_{ik} \star Y_{kj} + (-1)^{|X|} X_{ik} \star Q(Y)_{kj} + \Phi_{0ik} \star X_{kl} \star Y_{lj} - \\ &\quad - (-1)^{|X|+|Y|} X_{ik} \star Y_{kl} \star \Phi_{0lj} = (Q(X)_{il} + \Phi_{0ik} \star X_{kl} - (-1)^{|X|} X_{ik} \star \Phi_{0kl}) \star Y_{lj} + \\ &\quad + (-1)^{|X|} X_{ik} \star (Q(Y)_{kj} + \Phi_{0kl} \star Y_{lj} - (-1)^{|Y|} Y_{kl} \star \Phi_{0lj}) = \\ &\quad = \tilde{Q}(X)_{ik} \star Y_{kj} + (-1)^{|X|} X_{ik} \star \tilde{Q}(Y)_{kj} . \end{aligned} \quad (2.13)$$

²We do not write summation index for the summation over k, l in following expressions.

Finally we have

$$\begin{aligned}
\sum_{i,j} \int \tilde{Q}(X)_{ij} &= \sum_{i,j} \int Q(X)_{ij} + \int \Phi_{0ij} \star X_{ji} - (-1)^{|X|} \int X_{ij} \star \Phi_{0ji} = \\
&= \sum_{i,j} \int \Phi_{0ij} \star X_{ji} - (-1)^{|X|} (-1)^{|X||\Phi_0|} \int \Phi_{0ji} \star X_{ij} = 0 ,
\end{aligned} \tag{2.14}$$

using $|\Phi_0| = 1$. As a consequence the new BRST operator $\tilde{Q}$ obeys all axioms of the open bosonic string field theory even in the case when it is matrix valued. However it is desirable to have the BRST operator in the same form as the original one which means that will be diagonal in the CP indexes. In order to obtain such a form we should perform field redefinition according to [13]. We will see that with the solution given in (2.6) such a redefinition is very simple.

From (2.11) we immediately see that it is natural to perform field redefinition

$$\Psi_{ij} = e^{-\hat{K}}(\tilde{\Psi})_{ij} \tag{2.15}$$

that maps the new BRST operator (2.11) to the original one. It is important to stress that $\hat{K}$ is not arbitrary ghost number zero operator, while according to [10] $\hat{K}$ must obey following rules

$$\begin{aligned}
\hat{K}(\sum_l X_{il} \star Y_{lj}) &= \sum_l \hat{K}(X)_{il} \star Y_{lj} + \sum_l X_{il} \star \hat{K}(Y)_{lj} , \\
\sum_{i,j} \int \hat{K}(X_{ij} \star Y_{ji}) &= 0 .
\end{aligned} \tag{2.16}$$

If we presume that $\hat{K}$ obeys (2.16) it is a simple task to show that the field redefinition (2.15) leads to the action for the field $\tilde{\Psi}_{ij}$ that has the same form as (2.1) and where the BRST operator is the same as the original one Q . To prove this, let us start with the quadratic term

$$\begin{aligned}
\frac{1}{2} \sum_{i,j} \int \Psi_{ij} \star \tilde{Q}(\Psi_{ji}) &= \frac{1}{2} \sum_{i,j} \int e^{-\hat{K}}(\tilde{\Psi})_{ij} \star \tilde{Q}(e^{-\hat{K}}(\tilde{\Psi}))_{ji} = \\
&= \frac{1}{2} \sum_{i,j} \int \tilde{\Psi}_{ij} \star e^{\hat{K}}(\tilde{Q}(e^{-\hat{K}}(\tilde{\Psi})))_{ji} = \frac{1}{2} \sum_{i,j} \int \tilde{\Psi}_{ij} \star Q(\tilde{\Psi})_{ji} ,
\end{aligned} \tag{2.17}$$

where we have used

$$\begin{aligned}
\sum_{i,j} \int e^{-\hat{K}}(X)_{ij} \star Y_{ji} &= \sum_{i,j} \int X_{ij} \star Y_{ji} - \hat{K}(X)_{ij} \star Y_{ji} + \frac{1}{2} \hat{K}^2(X)_{ij} \star Y_{ji} + \dots = \\
&= \sum_{i,j} \int X_{ij} \star Y_{ji} + X_{ij} \star \hat{K}(Y)_{ji} - \frac{1}{2} \hat{K}(X)_{ij} \star \hat{K}(Y)_{ji} + \dots =
\end{aligned}$$

$$= \sum_{i,j} \int X_{ij} \star Y_{ji} + X_{ij} \star \hat{K}(Y)_{ji} + \frac{1}{2} X_{ij} \star \hat{K}^2(Y)_{ji} + \dots = \sum_{i,j} \int X_{ij} \star e^{\hat{K}}(Y)_{ji} , \quad (2.18)$$

and

$$e^{\hat{K}}(X \star Y)_{ij} = \sum_k e^{\hat{K}}(X)_{ik} \star e^{\hat{K}}(Y)_{kj} \quad (2.19)$$

that follows from (2.16). Now it is a simple task to show the invariance of the cubic term in (2.1) under field redefinition since

$$\begin{aligned} \sum_{i,j,k} \int \Psi_{ij} \star \Psi_{jk} \star \Psi_{ki} &= \sum_{i,j,k} \int e^{-\hat{K}}(\tilde{\Psi})_{ij} \star e^{-\hat{K}}(\tilde{\Psi})_{jk} \star e^{-\hat{K}}(\tilde{\Psi})_{ki} = \\ &= \sum_{i,j} \int \tilde{\Psi}_{ij} \star e^{\hat{K}}(e^{-\hat{K}}(\tilde{\Psi} \star \tilde{\Psi}))_{ji} = \sum_{i,j,k} \int \tilde{\Psi}_{ij} \star \tilde{\Psi}_{jk} \star \tilde{\Psi}_{ki} . \end{aligned} \quad (2.20)$$

From (2.17) and (2.20) we see that the action for the new fluctuation field $\tilde{\Psi}$ has the same form as the original one (2.1) which we wanted to prove.

Now from this general discussion we turn to the nonabelian extension of our result given in [14]. For that reason we must determine the form of the operator $\hat{K}$ and the meaning of the function $F_{ij}^I(\sigma)$ that appears in this operator. In fact, we introduce this function into $\hat{K}$ according to the analysis performed in [5, 7] in order to eliminate presence of the delta functions and also to eliminate contribution from the midpoint of the string. We demand that this function must obey these properties

$$F_{ij}^I(0) = F_{ij}^I(\pi) = \frac{1}{\sqrt{2\pi\alpha'}} Y_{ij}^I, F_{ij}^I\left(\frac{\pi}{2}\right) = 0 . \quad (2.21)$$

where Y_{ij}^I , $i, j = 1, \dots, N, I = 1, \dots, 25$ are matrices that describe the marginal deformation of the configuration of N D0-branes originally sitting in the origin. We also demand that these matrices commute among themselves so that can be put into the form

$$Y_{ij}^I = y_i^I \delta_{ij} , \quad (2.22)$$

where y_i^I parametrises shift of the position of the i -th D0-brane in the I -th direction.

Let us start with an infinitesimal form of the solution (2.6) Φ_{0ij} where we presume that the deformation parameters in Y_{ij}^I are small. Then we obtain

$$\Phi_{0ij} = Q(\hat{K}_L(\hat{I}, F))_{ij} . \quad (2.23)$$

Let us define operator $\hat{K}$ as follows ³

$$\begin{aligned} \hat{K}(X)_{ij} &= \sum_k (D_{ik} \star X_{kj} - X_{ik} \star D_{kj}) , D_{ik} = P_I^L(\mathcal{I}F^I)_{ik} , \\ P_I^L(F^I)_{ij} &= \frac{i}{2\pi\alpha'} \int_0^{\pi/2} d\sigma \eta_{IJ} \dot{X}^J F^I(\sigma)_{ij} , P_I^R(F^I) = \frac{i}{2\pi\alpha'} \int_{\pi/2}^{\pi} d\sigma \eta_{IJ} \dot{X}^J F^I(\sigma)_{ij} , \end{aligned} \quad (2.24)$$

³In the following we will not write its dependence of the function F_{ij}^I .

where in the definition of D_{ij} the summation over I is implicitly presumed. From previous expressions it is also clear the meaning of the symbols L, R that will appear in various operators. It is clear that $\hat{K}$ is nonabelian generalization of the operator K given in [14] for system of N D0-branes. We also see that $\hat{K}$ annihilates identity field $\hat{I} = \mathcal{I} \otimes \delta_{ij}$ since we have

$$\hat{K}(\hat{I})_{ij} = \sum_k D_{ik} \star \delta_{kj} \mathcal{I} - \delta_{ik} \mathcal{I} \star D_{kj} = D_{ij} - D_{ij} = 0 . \quad (2.25)$$

Then we define

$$\hat{K}_L(\hat{I})_{ij} = \sum_k D_{ik} \star \delta_{kj} \mathcal{I} = D_{ij} . \quad (2.26)$$

We can easily show that $\hat{K}$ obeys all axioms given in (2.16) since we have

$$\begin{aligned} \hat{K}(\sum_l X_{il} \star Y_{lj}) &= \sum_{k,l} D_{ik} \star X_{kl} \star Y_{lj} - X_{ik} \star Y_{kl} \star D_{lj} = \\ &= \sum_{k,l} D_{ik} \star X_{kl} \star Y_{lj} - X_{ik} \star D_{kl} \star Y_{lj} + \\ &\quad + X_{ik} \star D_{kl} \star Y_{lj} - X_{ik} \star Y_{kl} \star D_{lj} = \\ &= \sum_k \hat{K}(X)_{ik} \star Y_{kj} + X_{ik} \star \hat{K}(Y)_{kj} \end{aligned} \quad (2.27)$$

and

$$\begin{aligned} \text{Tr} \int \hat{K}(\Phi) &= \sum_{i,j} \left(\int D_{ij} \star \Phi_{ji} - \int \Phi_{ij} \star D_{ji} \right) = \\ &= \sum_{i,j} \left(\int D_{ij} \star \Phi_{ji} - (-1)^{|D||\Phi|} \int D_{ij} \star \Phi_{ji} \right) = 0 \end{aligned} \quad (2.28)$$

since the ghost number of D is equal to zero $|D| = 0$.

Now we return to (2.23) that can be written as [13, 14]

$$\Phi_{0ij} = Q(\hat{K}_L(\hat{I}))_{ij} = Q(P_I^L(\mathcal{I}F^I))_{ij} = [Q, P_I(F^I)]^L(\mathcal{I})_{ij} \quad (2.29)$$

and hence the new BRST operator is

$$\begin{aligned} \tilde{Q}(X)_{ij} &= \sum_k \delta_{ik} Q(X)_{kj} + \Phi_{0ik} \star X_{kj} - (-1)^{|X|} X_{ik} \star \Phi_{0kj} = \\ &= \sum_k \delta_{ik} Q(X)_{kj} - \tilde{D}^L(X)_{ij} - \tilde{D}^R(X)_{ij} \end{aligned} \quad (2.30)$$

using

$$\begin{aligned}
\sum_k \Phi_{0ik} \star X_{kj} &= \sum_k [Q, P_I(F_{ik}^I)]^L(\mathcal{I}) \star (X_{kj}) = - \sum_k \mathcal{I} \star [Q, P_I(F_{ik}^I)]^R(X_{kj}) , \\
&- \sum_k (-1)^{|X|} X_{ik} \star \Phi_{0kj} = -(-1)^{|X|} \sum_k X_{ik} \star [Q, P_I(F_{kj}^I)]^L(\mathcal{I}) = \\
&= (-1)^{|X|} \sum_k X_{ik} \star [Q, P_I(F_{kj}^I)]^R(\mathcal{I}) = - \sum_k [Q, P_I(F_{ik}^I)]^L(X_{kj})
\end{aligned} \tag{2.31}$$

and we also defined

$$\tilde{D}^L(X)_{ij} = \sum_k [Q, P_I(F_{ik}^I)]^L(X_{kj}), \quad \tilde{D}^R(X)_{ij} = \sum_k [Q, P_I(F_{ik}^I)]^R(X_{kj}) . \tag{2.32}$$

Firstly we must calculate $\tilde{D}^L$ which is equal to ⁴

$$\begin{aligned}
\tilde{D}_{ij}^L &= [Q^L, P_I^L(F^I)]_{ij} = \left[\frac{1}{\pi} \int_0^{\pi/2} d\sigma \left[c^0(\sigma) \frac{1}{2} ((2\pi\alpha')^2 P^2(\sigma) + X'^2(\sigma)) + \right. \right. \\
&\quad \left. \left. + c^1(\sigma) (2\pi\alpha') P_K X'^K(\sigma) \right], i \int_0^{\pi/2} d\sigma' P_I(\sigma') F^I(\sigma')_{ij} \right] = \\
&= \frac{i}{\pi} \int_0^{\pi/2} d\sigma \left(c^0(\sigma) \eta_{KL} X'^L(\sigma) + c^1(\sigma) (2\pi\alpha') P_K(\sigma) \right) \times \\
&\quad \times \partial_\sigma \int_0^{\pi/2} d\sigma' i \delta_D(\sigma, \sigma') \delta_I^K F^I(\sigma')_{ij} = \\
&= -\frac{1}{\pi} \int_0^{\pi/2} d\sigma \left(c^0(\sigma) X'^I(\sigma) \eta_{IJ} + c^1(\sigma) (2\pi\alpha') P_J(\sigma) \right) \partial_\sigma F^J(\sigma)_{ij} = \tilde{D}^L(F(\sigma)_{ij}) ,
\end{aligned} \tag{2.33}$$

where $\delta_D(\sigma, \sigma')$ is delta function with Dirichlet boundary conditions [14]. In order to determine the spectrum of the fluctuations around the new D0-brane configuration we perform field redefinition according to our general discussion. As it is clear from this construction the transformed BRST operator is the same as the original one

$$e^{\hat{K}} \left(\tilde{Q}(e^{-\hat{K}}) \right)_{ij} = Q \delta_{ij} \tag{2.34}$$

And the new field is equal to

$$\begin{aligned}
\tilde{\Psi} &= (\mathcal{I} + \hat{K})(\Psi)_{ij} = \Psi_{ij} + \sum_k P_I^L(\mathcal{I} F^I)_{ik} \star \Psi_{kj} - \Psi_{ik} \star P_I^L(\mathcal{I} F^I)_{kj} = \\
&= \Psi_{ij} - \sum_k P_I^R(F_{ik}^I \Psi_{kj}) - P_I^L(\Psi_{ik} F_{kj}^I) .
\end{aligned} \tag{2.35}$$

⁴In the following when we write Q inside the commutator we mean parts of Q whose commutators with X are nonzero, hence we ignore ghost contribution. More details can be found in [14].

Now we show that this string field carries correct winding charge that corresponds to the string joining i-Th and j-Th D0-brane. We know that the string winding charge is defined as

$$W^I = \int_0^\pi d\sigma \partial_\sigma X^I(\sigma) . \quad (2.36)$$

In order to see that this is conserved charge for the string obeying Dirichlet boundary conditions we can calculate its commutator with the bosonic Hamiltonian

$$H = \frac{1}{4\pi\alpha'} \int_0^\pi d\sigma ((2\pi\alpha')^2 P^2 + X'^2) \quad (2.37)$$

and we obtain

$$\begin{aligned} [W^I, H] &= \left[\int_0^\pi d\sigma X'^I(\sigma), \frac{1}{4\pi^2\alpha'} \int_0^\pi d\sigma' (2\pi\alpha')^2 P^2(\sigma') \right] \sim \\ &\sim \int_0^\pi d\sigma \int_0^\pi d\sigma' P^I(\sigma') \partial_\sigma \delta_D(\sigma', \sigma) \sim \int_0^\pi d\sigma \partial_\sigma \int_0^\pi d\sigma' P^I(\sigma') \delta_D(\sigma', \sigma) \sim \\ &\sim \int_0^\pi d\sigma \partial_\sigma P^I(\sigma) \sim P^I(\pi) - P^I(0) = 0 \end{aligned} \quad (2.38)$$

thanks to the Dirichlet boundary conditions on the boundary of the world-sheet. We also define

$$W^I = W_L^I + W_R^I, W_L^I = \int_0^{\pi/2} d\sigma \partial_\sigma X^I(\sigma), W_R^I = \int_{\pi/2}^\pi d\sigma \partial_\sigma X^I(\sigma), \quad (2.39)$$

so that we have

$$\begin{aligned} W^I(\tilde{\Psi}_{ij}) &= W^I(\Psi_{ij}) - W^I(P_J^R(\sum_k F_{ik}^J \Psi_{kj})) - W^I(P_J^L(\sum_k \Psi_{ik} F_{kj})) = \\ &= -[W_R^I, P_J^R(F^J)](\Psi)_{ij} - [W_L^I, P_J^L(F^J)](\Psi)_{ij} = \\ &= \sum_k (\Psi_{ik} Y_{kj}^I - Y_{ik}^I \Psi_{kj}) = \frac{1}{\sqrt{2\pi\alpha'}} (y_i^I - y_j^I) \Psi_{ij}, \end{aligned} \quad (2.40)$$

where we have used

$$\begin{aligned} W^I(P_J^L(F^J \Psi))_{ij} &= W_L^I(P_J^L(F^J \Psi))_{ij} + P_J^L(F^J W_R^I(\Psi))_{ij} = [W_L^I, P_J^L(F^J)](\Psi)_{ij}, \\ W^I \Psi_{ij} &= (W_L^I + W_R^I) \Psi_{ij} = 0 \Rightarrow W_L^I \Psi_{ij} = -W_R^I \Psi_{ij}, \end{aligned} \quad (2.41)$$

since the original configuration of D-branes corresponds to D-branes localized in the common point $x^I = 0$ so that the winding charge of the strings is equal to zero $W \Psi_{ij} = 0$. We also used

$$[W_L^I, P_J^L(F^J)_{ij}] = \left[\int_0^{\pi/2} d\sigma \partial_\sigma X^I(\sigma), i \int_0^{\pi/2} d\sigma' P_J(\sigma') F^J(\sigma')_{ij} \right] =$$

$$\begin{aligned}
&= \int_0^{\pi/2} d\sigma \partial_\sigma \int_0^{\pi/2} d\sigma' \delta_D(\sigma', \sigma) \delta_J^I F^J(\sigma')_{ij} = \\
&= \int_0^{\pi/2} d\sigma \partial_\sigma F^I(\sigma)_{ij} = (F^I(\pi/2) - F^I(0))_{ij} = -\frac{1}{\sqrt{2\pi\alpha'}} Y_{ij}^I, \\
&\quad [W_R^I, P_J^R(F^J)_{ij}] = (F^I(\pi) - F^I(\pi/2))_{ij} = \frac{1}{\sqrt{2\pi\alpha'}} Y_{ij}^I.
\end{aligned} \tag{2.42}$$

The previous result suggests that the new string field Ψ'_{ij} has a winding charge proportional to the distance between the i-Th and the j-Th D0-brane. For that reason the ground state $|ij\rangle$ of the string connecting i-Th and j-Th D0-brane can be represented with the operator

$$|ij\rangle = V_{ij} = \exp\left(i \int_0^\pi d\sigma P_I(\sigma) f_{ij}^I(\sigma)\right), \quad f^I(\sigma)_{ij} = \frac{1}{\sqrt{2\pi\alpha'}} \left(\frac{(y_i - y_j)^I}{\pi} \sigma - y_i^I\right) \tag{2.43}$$

and any redefined string field Ψ'_{ij} corresponding to the string connecting i-Th and j-Th D0-brane has a form

$$\Psi'_{ij} = \Psi_{ij} V_{ij}. \tag{2.44}$$

It is easy to see that (2.43) has correct winding charge. Firstly, we can calculate following commutator

$$\begin{aligned}
[W^I, V_{ij}] &= \left[W^I, \sum_{n=0}^{\infty} \frac{1}{n!} \left(i \int_0^\pi d\sigma' P_J(\sigma') f_{ij}^J(\sigma')\right)^N\right] = \\
&= \left[\int_0^\pi d\sigma \partial_\sigma X^I(\sigma), i \int_0^\pi d\sigma' P_J(\sigma') f_{ij}^J(\sigma') d\sigma'\right] V_{ij} = \\
&= - \int_0^\pi d\sigma \partial_\sigma \int_0^\pi d\sigma' \delta_J^I \delta_D(\sigma', \sigma) f_{ij}^J(\sigma') V_{ij} = - \int_0^\pi d\sigma \partial_\sigma f_{ij}^I(\sigma) V_{ij} = \\
&= -[f^I(\pi)_{ij} - f^I(0)_{ij}] V_{ij} = \frac{1}{\sqrt{2\pi\alpha'}} (y_j - y_i)^I V_{ij}.
\end{aligned} \tag{2.45}$$

We see that V_{ij} has the correct winding charge. Now we will calculate the action of original BRST operator Q on general new string field $\Psi'_{ij} = V_{ij} \Psi_{ij}$. First of all, it is clear that the action of Q on the string field Ψ is the same as in the original action. For our purposes it is important the action of Q on V_{ij} . For that reason let us calculate

$$\begin{aligned}
[X^I(\sigma), V_{ij}] &= \left[X^I(\sigma), \sum_{N=0}^{\infty} \frac{1}{N!} \left(i \int_0^\pi d\sigma' P_J(\sigma') f_{ij}^J(\sigma')\right)^N\right] = \\
&= \left[X^I(\sigma), i \int_0^\pi d\sigma' P_J(\sigma') f_{ij}^J(\sigma')\right] V_{ij} = - \int_0^\pi d\sigma' \delta_J^I \delta_D(\sigma, \sigma') f_{ij}^J(\sigma') V_{ij} = -f_{ij}^I(\sigma) V_{ij}
\end{aligned} \tag{2.46}$$

and consequently

$$\begin{aligned}
[Q, V_{ij}] &= \left[\frac{1}{\pi} \int_0^\pi d\sigma \left(\frac{1}{2} c^0(\sigma) \partial_\sigma X^I(\sigma) \partial_\sigma X^J(\sigma) \eta_{IJ} + \right. \right. \\
&\quad \left. \left. + c^1(\sigma) \partial_\tau X^I(\sigma) \partial_\sigma X^J(\sigma) \eta_{IJ} \right), V_{ij} \right] = \\
&= \frac{1}{\pi} \int_0^\pi d\sigma \frac{(y_j^I - y_i^I)}{\sqrt{2\pi\alpha'}} V_{ij} \eta_{IJ} \left(c^0(\sigma) \partial_\sigma X^J(\sigma) + c^1(\sigma) \partial_\tau X^J(\sigma) \right) + \\
&\quad + \frac{1}{\pi} \int_0^\pi d\sigma c^0(\sigma) \frac{(y_j^I - y_i^I) \eta_{IJ} (y_j^J - y_i^J)}{4\pi\alpha'} V_{ij} ,
\end{aligned} \tag{2.47}$$

where we have used

$$\begin{aligned}
&\left[\frac{1}{2\pi} \int_0^\pi d\sigma c^0(\sigma) \partial_\sigma X^I(\sigma) \partial_\sigma X^J(\sigma) \eta_{IJ}, V_{ij} \right] = \\
&= \frac{1}{2\pi} \int_0^\pi d\sigma c^0(\sigma) \left[2\partial_\sigma [X^I(\sigma), V_{ij}] \partial_\sigma X^J(\sigma) \eta_{IJ} + \right. \\
&\quad \left. + \partial_\sigma [X^I(\sigma), \partial_\sigma [X^J(\sigma), V_{ij}]] \eta_{IJ} \right] ;
\end{aligned} \tag{2.48}$$

$$\begin{aligned}
&\frac{1}{\pi} \int_0^\pi d\sigma c^0(\sigma) \partial_\sigma [X^I(\sigma), V_{ij}] \partial_\sigma X^J(\sigma) \eta_{IJ} = \\
&= \frac{1}{\pi} \int_0^\pi d\sigma c^0(\sigma) \partial_\sigma (-f_{ij}^I(\sigma)) V_{ij} \partial_\sigma X^J(\sigma) \eta_{IJ} = \\
&= \frac{1}{\pi} \int_0^\pi d\sigma c^0(\sigma) \frac{(y_j^I - y_i^I)}{\sqrt{2\pi\alpha'}} V_{ij} \eta_{IJ} ;
\end{aligned} \tag{2.49}$$

$$\begin{aligned}
&\frac{1}{2\pi} \int_0^\pi d\sigma c^0(\sigma) \partial_\sigma [X^I(\sigma), [X^J(\sigma), V_{ij}]] \delta_{IJ} = \\
&= \frac{1}{2\pi} \int_0^\pi d\sigma c^0(\sigma) \partial_\sigma [X^I(\sigma), \frac{(y_j^J - y_i^J)}{\sqrt{2\pi\alpha'}} V_{ij}] = \\
&= \frac{1}{2\pi} \int_0^\pi d\sigma c^0(\sigma) \frac{(y_j^I - y_i^I) \eta_{IJ} (y_j^J - y_i^J)}{4\pi\alpha'} V_{ij} ;
\end{aligned} \tag{2.50}$$

$$\begin{aligned}
&\left[\frac{1}{2\pi} \int_0^\pi d\sigma c^1(\sigma) \partial_\tau X^I(\sigma) \partial_\sigma X^J(\sigma), V_{ij} \right] = \\
&= \frac{1}{2\pi} \int_0^\pi d\sigma c^1(\sigma) \partial_\tau X^I(\sigma) \eta_{IJ} \partial_\sigma (-f(\sigma)_{ij}^J V_{ij}) = \\
&= \frac{1}{2\pi} \int_0^\pi d\sigma c^1(\sigma) \partial_\tau X^I(\sigma) \eta_{IJ} \frac{(y_j^J - y_i^J)}{\sqrt{2\pi\alpha'}} V_{ij} .
\end{aligned} \tag{2.51}$$

We see that (2.47) describes the right action of the BRST operator Q on the string state corresponding to the winding mode. Hence we can claim that the solution given in (2.23) correctly describes small marginal deformation in case of the system of N D0-branes. We also see that using of the function F_{ij}^I leads to the completely non singular solution with accord with [7].

3. Finite deformation

As in the abelian case [14] we can easily extend this approach to the case of finite deformation when the solution (2.6) can be explicitly written as

$$\Phi_{0ij} = Q(\hat{I})_{ij} + Q(\hat{K}_L(\hat{I}))_{ij} + \sum_{n=2}^{\infty} \frac{1}{n!} [[Q(\hat{K}_L(\hat{I}), \hat{K}_L(\hat{I})), \hat{K}_L(\hat{I})], \dots]_{ij}, \quad (3.1)$$

where

$$[Q(\hat{K}_L(\hat{I}), \hat{K}_L(\hat{I}))]_{ij} = \sum_k Q(\hat{K}_L(\hat{I}))_{ik} \star \hat{K}_L(\hat{I})_{kj} - \hat{K}_L(\hat{I})_{ik} \star Q(\hat{K}_L(\hat{I}))_{kj} \quad (3.2)$$

and commutators of higher orders are defined in the same manner. Now we will proceed in the same way as in [14]. The first term in (3.1) is given in (2.29), (2.33). The second term is equal to

$$\begin{aligned} \frac{1}{2} [Q(\hat{K}_L(\hat{I}), \hat{K}_L(\hat{I}))]_{ij} &= \frac{1}{2} \sum_k Q(\hat{K}_L(\hat{I}))_{ik} \star \hat{K}_L(\hat{I})_{kj} - \hat{K}_L(\hat{I})_{ik} \star Q(\hat{K}_L(\hat{I}))_{kj} = \\ &= \frac{1}{2} \sum_k \tilde{D}^L(\mathcal{I})_{ik} \star P_J^L(\mathcal{I}F^J)_{kj} - P_J^L(\mathcal{I}F^J)_{ik} \star \tilde{D}^L(\mathcal{I})_{kj} = \\ &= \frac{1}{2} \sum_k P_J^L(\tilde{D}^L(\mathcal{I})_{ik} F_{kj}^J) - \tilde{D}^L(P_J(\mathcal{I}F^J)_{ik} F(\sigma)_{kj}), \end{aligned} \quad (3.3)$$

where

$$\begin{aligned} \sum_k P_K^L(\tilde{D}^L(\mathcal{I})_{ik} F_{kj}^K) &= \sum_k i \int_0^{\pi/2} d\sigma' P_K(\sigma') \left(-\frac{1}{\pi} \int_0^{\pi/2} d\sigma (c^0(\sigma) \partial_\sigma X^I(\sigma) \eta_{IJ} + \right. \\ &\quad \left. + c^1(\sigma) (2\pi\alpha') P_J(\sigma)) \partial_\sigma F^J(\sigma)_{ik} \right) F_{kj}^K(\sigma') = \sum_k \\ &= \left(-\frac{1}{\pi} \int_0^{\pi/2} d\sigma (c^0(\sigma) \partial_\sigma X^I(\sigma) \eta_{IJ} + c^1(\sigma) (2\pi\alpha') P_J(\sigma)) \right) \times \\ &\quad \times i \int_0^{\pi/2} d\sigma' P_K(\sigma') F^K(\sigma')_{ik} \partial_\sigma F_{kj}^J(\sigma) - \\ &\quad - \frac{i}{\pi} \int_0^{\pi/2} d\sigma' \int_0^{\pi/2} d\sigma c^0(\sigma) \partial_\sigma (-i\delta_D(\sigma, \sigma')) \delta_K^I \eta_{IJ} \partial F_{ik}^J(\sigma) F_{kj}^K(\sigma') = \\ &= \sum_k \tilde{D}^L(P_K(\mathcal{I}F^K)_{ik} F_{kj}(\sigma)) - \frac{1}{\pi} \int_0^{\pi/2} d\sigma c^0(\sigma) \partial_\sigma F_{ik}^I(\sigma) \eta_{IJ} \partial_\sigma \int_0^{\pi/2} d\sigma' \delta_D(\sigma, \sigma') F_{kj}^J(\sigma'). \end{aligned} \quad (3.4)$$

Then (3.3) is equal to

$$\frac{1}{2} [Q(\hat{K}_L(\hat{I}), \hat{K}_L(\hat{I}))] = -\frac{1}{2\pi} \int_0^{\pi/2} d\sigma c^0(\sigma) \partial_\sigma F_{ik}^I(\sigma) \eta_{IJ} \partial_\sigma F_{kj}^J(\sigma) \quad (3.5)$$

and according to this result we immediately see that the higher order commutators between $\hat{K}$ in (3.1) are equal to zero and we obtain an exact result

$$\begin{aligned} \Phi_{0ij} = & -\frac{1}{\pi} \int_0^{\pi/2} d\sigma \left(c^0(\sigma) X'_I(\sigma) \eta_{IJ} + c^1(\sigma) (2\pi\alpha') P_J(\sigma) \right) \partial_\sigma F^J(\sigma)_{ij} - \\ & -\frac{1}{2\pi} \sum_k \int_0^{\pi/2} d\sigma c^0(\sigma) \partial_\sigma F^I_{ik}(\sigma) \eta_{IJ} \partial_\sigma F^J_{kj}(\sigma) . \end{aligned} \quad (3.6)$$

As in previous section we perform field redefinition, following general analysis given in [13]. Then the new BRST operator $\tilde{Q}_{ij}$ is mapped to the original one $Q\delta_{ij}$ and the new string field $\tilde{\Psi}_{ij}$ is equal to

$$\tilde{\Psi}_{ij} = \exp(\hat{K})(\Psi)_{ij} = \sum_{n=0}^{\infty} \frac{1}{n!} \hat{K}^n(\Psi)_{ij} , \quad (3.7)$$

where we obtain from (2.35)

$$\hat{K}(\Psi)_{ij} = \sum_k P_I^L(\mathcal{I}F^I)_{ik} \star \Psi_{kj} - \Psi_{ik} \star P_I^L(\mathcal{I}F^I)_{kj} = -\sum_k P_I^R(F^I_{ik} \Psi_{kj}) - P_I^L(\Psi_{ik} F^I_{kj}) , \quad (3.8)$$

and from (2.40)

$$[W^I, \hat{K}](\Psi)_{ij} = (y_i^I - y_j^I) \Psi_{ij} . \quad (3.9)$$

As in the case of the small deformation let us work out the action of W on $\tilde{\Psi}$. We obtain

$$W(\tilde{\Psi})_{ij} = W(e^{\hat{K}}(\Psi))_{ij} = [W, e^{\hat{K}}](\Psi)_{ij} = (y_i^I - y_j^I) \tilde{\Psi}_{ij} , \quad (3.10)$$

where we have again used the fact that $W(\Psi)_{ij} = 0$ and also

$$[W, e^{\hat{K}}] = \left[W, \sum_{N=0}^{\infty} \frac{1}{N!} \hat{K}^N \right] = \sum_{N=0}^{\infty} \frac{1}{N!} \hat{K}^N ([W, \hat{K}]) = e^{\hat{K}}([W, \hat{K}]) . \quad (3.11)$$

It is important to stress that the upper formula holds only on condition $[[W, \hat{K}], \hat{K}] = 0$.

We immediately see that the new string field $\tilde{\Psi}_{ij}$ has the correct winding number proportional to the distance between i-Th and j-Th D0-branes. As in the previous section we can write this field using the winding state V_{ij} and perform the same calculation. As a result, the action for the fluctuation field has the same form as the original one

$$S = -\frac{1}{2} \sum_{i,j} \int \tilde{\Psi}_{ij} \star Q(\tilde{\Psi}_{ji}) - \frac{1}{3} \sum_{i,j,k} \int \tilde{\Psi}_{ij} \star \tilde{\Psi}_{jk} \star \tilde{\Psi}_{ki} . \quad (3.12)$$

In this section we found the exact string field theory solution (3.6) describing marginal deformation of the configuration of N D0-branes from the background corresponding to N coincided D0-branes to the background where D0-branes are localized in general positions. We believe that our result could be helpful for other study marginal solutions in the string field theory.

4. Conclusion

In this paper we have extended our solutions of the open bosonic string field theory that were found in [14] to the case of the open bosonic string field theory defined on the background of N D0-branes. We have shown that there exist exact solution of the open bosonic string field theory that describes any marginal deformation of the original configuration of N D0-branes localized in the origin of the space to the background configuration of N D0-branes in general positions. We have also seen that using the regulation of the solution, following [5, 6, 7] we have obtained solutions that is completely non singular. This fact can be used as an explanation of the presence of the singular term in the exact solution given in our previous work [14].

The extension of this work is obvious. First of all, we would like to perform similar analysis in the exact conformal field theory language, following seminal papers [10, 11, 12]. We would like to study other solutions using our formalism. We hope to return to these questions in the future.

Acknowledgment

We would like to thank Ulf Danielsson and Ulf Lindstrom for their support in our work. This work is partly supported by EU contract HPRN-CT-2000-001222. This work is also supported by the Czech Ministry of Education under Contract No 143100006.

References

- [1] K. Ohmori, "*A Review on Tachyon Condensation in Open String Field Theories*," [hep-th/0102085](#).
- [2] P. J. De Smet, "*Tachyon Condensation: Calculation in String Field Theory*," [hep-th/0109182](#).
- [3] I. Ya Arefeva, D. M. Belov, A. A. Girgavets, A. S. Koshelev and P. B. Medvedev, "*Non-commutative Field Theories and (Super)string Field Theories*," [hep-th/0111208](#).
- [4] E. Witten, "*Noncommutative Geometry and String Field Theory*," *Nucl. Phys.* **B 268** (1986) 253.
- [5] T. Takahashi and S. Tanimoto, "*Wilson Lines and Classical Solutions in Cubic Open String Field Theory*," *Prog. Theor. Phys.* **‘06** (2001) 863, [hep-th/0107046](#).
- [6] T. Takahashi and S. Tanimoto, "*Dilaton Condensation in Cubic Open String Field Theory*," [hep-th/0112124](#).
- [7] T. Takahashi and S. Tanimoto, "*Marginal and scalar solutions in cubic open string field theory*," [hep-th/0202133](#).

- [8] A.Sen and B. Zwiebach, “*Large Marginal Deformation in String Field Theory,*” *J. High Energy Phys.* **0010** (2000) 009, [hep-th/0007153](#).
- [9] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, “*Patterns in open string field theory solutions,*” *J. High Energy Phys.* **0203** (2002) 003, [hep-th/0201159](#).
- [10] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, “*Ghost Structure and Closed Strings in Vacuum String Field Theory,*” [hep-th/0111129](#).
- [11] I. Kishimoto and K. Ohmori, “*CFT Description of Identity String Field: Toward Derivation of the VSFT Action,*” [hep-th/0112169](#).
- [12] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, “*Star algebra projectors,*” [hep-th/0202151](#).
- [13] J. Klusoň, “*Some Solutions of Berkovits’ Superstring Field Theory,*” [hep-th/0201054](#).
- [14] J. Klusoň, “*Exact Solutions of Open Bosonic String Field Theory,*” [hep-th/0202045](#).
- [15] J. Klusoň, “*Matrix model and string field theory,*” [hep-th/0011029](#).